

Table 4E
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by Region
Children Aged < 5 Years ⁽²⁾

Page: 1
Run Date: 3 / 13 / 09

Comparison of Racial and Ethnic Distribution by Region								
Region	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
NERO	24,623	56.3	2.9	23.9	3.8	7.1	6.0	0.0
NRO	10,675	69.9	0.8	8.0	7.1	8.7	5.6	0.0
SERO	56,212	20.9	37.3	33.9	0.3	2.9	4.7	0.1
SRO	20,429	57.8	10.0	24.1	0.4	2.9	4.8	0.0
WRO	17,709	77.0	1.4	8.9	1.8	4.6	6.3	0.0
Wisconsin	129,648	45.1	18.5	24.9	1.7	4.4	5.2	0.0
Nation (Prior Year)	7,995,600	31.6	19.4	40.9	1.0	2.5	1.8	2.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5E
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by Region
Children Aged < 5 Years ⁽²⁾

Page: 1
Run Date: 3 / 13 / 09

Comparison of Age Distribution by Region				
Region	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
NERO	24,623	33.6	21.7	44.8
NRO	10,675	35.3	20.3	44.4
SERO	56,212	35.4	21.4	43.2
SRO	20,429	36.3	20.9	42.7
WRO	17,709	34.5	20.8	44.8
Wisconsin	129,648	35.0	21.2	43.8
Nation (Prior Year)	7,995,600	37.6	22.1	40.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 6E
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Region
Children Aged < 5 Years (2)

Page: 1
Run Date: 3 / 13 / 09

Comparison of Growth and Anemia Indicators by Region (3,4)													
Region	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Obese (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
NERO	5,938	8.4 (4)	8.1 (3)	24,069	5.1 (2)	3.9 (5)	13.5 (4)	10,881	15.7 (2)	13.6 (3)		17,539	7.2 (2)
NRO	2,510	7.7 (3)	8.0 (2)	10,462	5.9 (4)	2.8 (1)	13.1 (2)	4,658	16.6 (4)	13.9 (4)		7,700	6.1 (1)
SERO	14,213	9.4 (5)	6.4 (1)	54,899	5.9 (4)	3.6 (4)	13.5 (4)	23,885	16.3 (3)	13.9 (4)		39,312	15.2 (5)
SRO	5,114	7.5 (2)	8.7 (4)	19,958	4.9 (1)	3.2 (3)	13.1 (2)	8,612	15.1 (1)	12.9 (1)		14,348	8.4 (3)
WRO	4,165	7.0 (1)	9.5 (5)	17,381	5.4 (3)	3.0 (2)	12.4 (1)	7,839	17.9 (5)	13.1 (2)		12,720	9.2 (4)
Wisconsin	31,940	8.5	7.6	126,769	5.5	3.5	13.3	55,875	16.3	13.6		91,619	11.0
Nation (Prior Year)	1,953,242	9.1	6.4	7,526,585	6.2	4.5	13.9	3,058,130	16.4	14.9		4,759,588	15.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this region's rate to other regions. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7E
2008 Pediatric Nutrition Surveillance (1)
Wisconsin

Page: 1
Run Date: 3 / 13 / 09

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Region (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Region
(4,5)

	Breastfeeding (5)						Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household	
Region	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
NERO	5,940	62.4 (4)	2,642	25.2 (3)	3,883	10.2 (5)	1,952	16.0 (4)	2,469	7.0 (4)	11,705	81.6 (4)	24,312	8.8 (3)
NRO	2,513	61.4 (5)	1,167	24.9 (5)	1,507	11.3 (4)	828	19.0 (2)	1,051	8.6 (3)	5,049	84.8 (2)	10,565	9.4 (4)
SERO	14,232	63.2 (3)	5,914	25.0 (4)	8,753	11.8 (3)	4,122	11.4 (5)	5,565	3.5 (5)	25,892	79.7 (5)	55,377	10.5 (5)
SRO	5,124	68.0 (2)	2,347	34.2 (1)	3,136	16.7 (1)	1,851	18.8 (3)	2,155	8.8 (2)	9,325	83.1 (3)	20,089	8.2 (2)
WRO	4,176	70.0 (1)	1,893	28.9 (2)	2,583	12.2 (2)	1,407	22.1 (1)	1,723	10.0 (1)	8,448	85.2 (1)	17,607	7.1 (1)
Wisconsin	31,985	64.6	13,963	27.1	19,862	12.3	10,160	15.7	12,963	6.3	60,419	81.8	127,950	9.3
Nation (Prior Year)	1,747,004	59.8	874,434	25.4	979,105	17.5	127,047	10.2	182,064	5.2	687,613	75.0	3,047,308	11.3

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this region's rate to other regions. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6E
2006 - 2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Region
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Region (3,4)													
Region	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)			Under- weight (7)	Obese (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
NERO	17,501	8.3 (4)	7.9 (2)	69,642	5.1 (1)	4.0 (5)	12.7 (3)	31,291	15.8 (2)	13.2 (4)		50,471	7.3 (2)
NRO	7,469	7.1 (1)	8.5 (3)	30,048	6.0 (5)	3.0 (1)	12.4 (2)	13,175	16.8 (4)	12.9 (2)		21,751	5.5 (1)
SERO	41,907	9.5 (5)	6.3 (1)	158,514	5.8 (4)	3.9 (4)	12.8 (4)	68,095	16.3 (3)	13.6 (5)		111,841	15.2 (5)
SRO	14,901	8.1 (3)	8.8 (4)	56,824	5.3 (2)	3.2 (2)	12.8 (4)	24,217	15.5 (1)	13.0 (3)		40,224	8.7 (4)
WRO	12,363	7.2 (2)	9.2 (5)	50,320	5.4 (3)	3.3 (3)	12.0 (1)	22,379	17.2 (5)	12.7 (1)		36,350	8.5 (3)
Unknown Regions	0	* *	* *	40	* *	* *	* *	21	* *	* *		33	* *
Wisconsin	94,141	8.6	7.6	365,388	5.5	3.6	12.6	159,178	16.2	13.3		260,670	10.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this region's rate to other regions. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7E
2006 - 2008 Pediatric Nutrition Surveillance (1)
Wisconsin

Page: 1
Run Date: 3 / 13 / 09

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Region (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Region
(4,5)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Region (4,5)														
	Breastfeeding (5)						Exclusive Breastfeeding (6)			≤2 Hours TV Viewing/Day (7)		Smoking in Household		
Region	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
NERO	18,892	62.4 (3)	12,401	20.7 (4)	12,615	10.8 (5)	11,253	10.7 (4)	11,058	5.1 (4)	33,762	81.0 (4)	39,433	10.4 (2)
NRO	8,060	61.1 (5)	5,414	20.7 (4)	5,152	11.5 (3)	4,788	13.3 (3)	4,655	6.5 (3)	14,329	83.4 (3)	16,864	10.8 (3)
SERO	44,557	62.3 (4)	28,745	21.8 (3)	28,883	11.9 (2)	25,789	7.8 (5)	25,573	3.5 (5)	73,678	78.6 (5)	89,058	12.8 (5)
SRO	16,215	68.1 (2)	10,676	29.1 (1)	10,274	16.2 (1)	9,423	14.2 (2)	9,113	7.7 (1)	26,131	84.4 (2)	32,697	10.9 (4)
WRO	13,381	70.0 (1)	8,947	23.3 (2)	8,785	11.2 (4)	8,025	14.5 (1)	7,811	6.9 (2)	24,120	84.9 (1)	28,224	8.4 (1)
Unknown Regions	0	* *	2	* *	2	* *	0	* *	2	* *	21	* *	30	* *
Wisconsin	101,105	64.2	66,185	22.9	65,711	12.2	59,278	10.7	58,212	5.1	172,041	81.3	206,306	11.3

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this region's rate to other regions. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.